

Guaranteed composition Nitrogen (N) 20%, phosphorus (as P₂O₅) 20%, potassium (as K₂O) 20%, magnesium (Mg) 0.9%, boron (B) 0.0315, cobalt (Co) 0.0012, copper (Cu) 0.0750, iron (Fe) 0.2420, manganese (Mn) 0.0750, molybdenum (Mo) 0.0012, zinc (Zn) 0.0750 % w/v TpTep is a 20:20:20 high concentration suspension fertiliser with extra elements and microelements for foliar applications. TpTep can be used on a range of indoor and outdoor crops with a requirement for a full complement of nutrients

TIPTOP has been specially formulated to ensure it is fully stable in the bottle and fully water soluble, ensuring consistent rates of application and dosing. The TIPTOP formulation has been designed to give an optimal spray pattern, ensuring thorough coverage of the plant giving a more even spread of nutrients.

The full compliment of macro and micro-elements ensure that TIPTOP can be used as a foliar feed without the necessity for base dressings. Containing 'Lightning' technology, the nutrients are rapidly taken up by the plant addressing nutrient deficiencies quickly and effectively with minimal losses.

Recommendations: Cereals, OSR, Top fruit trees, potato, all vegetables, all fruit, tomato, all greenhouse crops and most other crops.

| Crop | Application | Growth Stage |
|---------------------------|----------------------|----------------------------------|
| OSR | Foliar | After Drilling to end of pod set |
| Cereals | Foliar | From Planting Date |
| Sugar beet | Foliar | From Planting Date |
| Potatoes | Foliar | From Planting Date |
| Pulses | Foliar | From Planting Date |
| Vegetables | Foliar | From Planting Date |
| Protected Crops | Irrigation or foliar | From Planting Date |
| Tomato | Irrigation or foliar | From Planting Date |
| Strawberry and soft fruit | Irrigation or foliar | From Planting Date |
| Floriculture | Irrigation or foliar | From Planting Date |
| Top fruit and vineyards | | |
| Citrus | Irrigation or foliar | |
| Vineyards and grapes | Irrigation or foliar | |

Application Rate

TipTop can be used as at a rate of 2-5 litres / Ha for most crops,

depending on the requirement for feed. Use in a minimum of 200 litres of water/Ha.

For indoor crops or foliar sprays use 1-2 ml / litre water and ensure thorough coverage, apply at 400 - 1000 litres / ha.

Repeat applications as necessary to promote vigorous, healthy growth.



Causes severe skin burns and eye damage.

Precautionary statements:

Do not breathe dust/fumes/gas/mist/vapours/spray.

Wear protective gloves/protective clothing/eye protection/face protection.

IF SWALLOWED: rinse mouth. Do NOT induce vomiting.

IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water/shower.

IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.



Tel: 01673 838 143 Email: info@bionatureagriculture.com www.bionatureagriculture.com Caistor Lane, Tealby, Market Rasen, Lincolnshire, LN8 3XW TipTop is a Trade Mark of Ecoculture Biosciences TipTop contains Ecoculture Biosciences Technology



